

**R13**

Code No: 113AA

JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD

B.Tech II Year I Semester Examinations, November - 2015

MATHEMATICS – II

(Common to CE, CHEM, MMT, AE, PTE, CEE)

Time: 3 Hours

Max. Marks: 75

**Note:** This question paper contains two parts A and B.

Part A is compulsory which carries 25 marks. Answer all questions in Part A.

Part B consists of 5 Units. Answer any one full question from each unit.

Each question carries 10 marks and may have a, b, c as sub questions.

**PART- A****(25 Marks)**

- 1.a) If  $\phi$  satisfies Laplacian equation, show that  $\Delta\phi$  is both solenoidal. [2M]
- b) Find curl  $x^2i + yj + zk$ . [3M]
- c) Write the Dirichlet's conditions for the existence of Fourier series of a function  $f(x)$  in the interval  $(\alpha, \alpha + 2\pi)$ . [2M]
- d) State and prove linearity property of Fourier transforms. [3M]
- e) State interpolation. [2M]
- f) If the interval of difference is unity, Find  $\Delta(x^2 + 2x)$ . [3M]
- g) Under what conditions Gauss-Seidal method is applicable. [2M]
- h) Derive an iteration formula to find square root of a number n by Newton Raphson method. [3M]
- i) State merits and demerits of Runge-Kutta method. [2M]
- j)

x	0	1	2	3
y	1	0.5	0.3333	0.25

Evaluate  $\int_0^3 \frac{dx}{1+x}$  by Trapezoidal rule.

[3M]

**PART-B****(50 Marks)**

2. State and verify Stokes theorem for the function  $\vec{f} = x^2i + xyj$  integrated round the square in the plane  $z = 0$  whose sides are along the lines  $x = 0 = y$ ,  $x = a = y$ . [10]

**OR**

3. Find the work done in moving in a particle in the force field  $\vec{f} = 3x^2i + (2zx - y)j + zk$ , along the curve defined by  $x^2 = 4y$ ,  $3x^3 = 8z$  from  $x = 0$  to  $x = 2$ . [10]

- 4.a) Expand the Fourier series expansion of the function

$$f(x) = 0, -\pi \leq x \leq 0$$

$$= \sin x, 0 \leq x \leq \pi$$

$$\text{and deduce } \frac{1}{1.3} - \frac{1}{3.5} + \frac{1}{5.7} - \dots = \frac{\pi - 2}{4}$$

- b) Obtain the half range cosine series for the function  $f(x) = x^2$  when  $0 < x < \pi$  and

$$\text{find the sum of the series } \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots$$

[5+5]

OR

- 5.a) State and prove Change of scale property of Fourier transforms.  
b) Find Fourier sine transform of  $f(x) = \frac{1}{x(x^2 + a^2)}$  and hence deduce cosine transform of  $\frac{1}{x^2 + a^2}$ . [5+5]

- 6.a) Find the missing term in the following data:

x	0	1	2	3	4
y	1	3	9	-	81

- Why this value is not equal to  $3^3$ . Explain.  
b) Fit a straight line to the following data by the method of least square. [5+5]

x	0	1	2	3	4
y	1	1.8	3.3	4.5	6.5

OR

- 7.a) Given  $\sin 45^\circ = 0.7071$ ,  $\sin 50^\circ = 0.7660$ ,  $\sin 55^\circ = 0.8192$  and  $\sin 60^\circ = 0.8660$ . Find  $\sin 52^\circ$  using Newton's interpolation formula.  
b) A curve passes through the points (0, 18), (1, 10), (3, -18) and (6, 90). Find the slope of the curve at  $x = 2$ . [5+5]

8. Find the inverse of the matrix  $\begin{bmatrix} 2 & 3 & 1 \\ 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix}$  using LU decomposition method. [10]

OR

- 9.a) Find the root of  $2x - \log x = 7$ , correct to three places of decimal using iteration method.  
b) Find the root of the equation  $x \log_{10}(x) = 1.2$  using False position method. [5+5]
10. Find the successive approximate solution of the differential equation  $y' = y$ ,  $y(0) = 1$  by Picard's method and compare it with exact solution. [10]

OR

11. Determine the largest Eigen value and the corresponding eigenvector of the matrix  $\begin{bmatrix} 4 & 1 & 0 \\ 1 & 20 & 1 \\ 0 & 1 & 4 \end{bmatrix}$  to 3 correct decimal places using the power method. [10]

---ooOoo---